

FOR DISTRIBUTION AND USE ONLY UNDER SECTION 18 EMERGENCY EXEMPTION
This labeling must be in the possession of the user at the time of pesticide application.

FujiMite[®] 5EC Miticide/Insecticide

(EPA Reg. No. 71711-19)

**FOR USE ON HOPS ONLY FOR THE CONTROL OF MITES
IN THE STATE OF IDAHO**

This specific exemption is effective from May 31, 2005 to September 15, 2005

EPA File Symbol 05 – ID – 11

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read entire Directions for Use and Disclaimer of Warranties before using this product.

HOPS

For best results, apply before mite populations exceed 5 mites per leaf. For treatments when there is less than a full canopy of growth, the total amount of FujiMite 5EC used per acre may be reduced according to the percentage of canopy sprayed compared to the normal full canopy. Do not use less than two pints FujiMite 5EC per acre on hops that have grown to full height.

RECOMMENDED APPLICATION RATES FOR FUJIMITE 5EC ON HOPS

Pest	Application Rate Per Acre	Specific Use Directions and Restrictions
Mites (such as Two-spotted spider mite, Willamette spider mite, Pacific spider mite, etc.)	2 to 3 pints (32 - 48 fl oz) (0.1 - 0.15 lb ai)	<ul style="list-style-type: none">• Use in sufficient volume to obtain uniform plant coverage using properly calibrated high or low volume spray equipment.• Spray concentrations above 100 ppm are recommended with full coverage high volume spray equipment.• Do not apply within 21 days of harvest.• A maximum of 2 ground applications at a rate of 32 - 48 fl oz product (0.1 - 0.15 lb ai) per acre may be made per season.• Do not exceed 96 fluid ounces of product per acre per season.• Do not apply this product by air or through any type of irrigation system. Applications may be made using ground equipment only.• Do not apply within 75 feet of all aquatic areas.• Do not make subsequent applications without rotating to at least two other miticide products between applications.

The following table is a guide to assist in calculating the appropriate spray concentration:

Amount of FujiMite 5EC for 100 ppm Spray Concentration	
Spray Volume per Acre	Pints FujiMite 5EC
75	1.2
100	1.6
125	2.0
150	2.4
188	3.0

Nichino America, Inc.

WARNING

Leaf yellowing may occur when FujiMite 5EC is combined with spray oil in excess of 1% of the spray volume. If this symptom occurs, it is usually more pronounced on newly expanding leaves. This symptom may occur in plants under stress and is worsened by certain conditions including, but not limited to, the following:

- High Temperatures (air temperatures exceeding 90°F at the time of application or within a few days after application)
- Wet Soil Conditions and High Humidity (rainy, misty, or foggy weather within a few days after application)
- Storm Damage (including hail and wind)

IMPORTANT

This label must be in the possession of the user at the time of application.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed. This product is sold subject to the Conditions of Sale set forth on the container label.

Any adverse effects resulting from the use of FujiMite 5EC under this emergency exemption must be immediately reported to the Idaho Department of Agriculture.

© 2005. Nichino America, Inc.
FujiMite® is a trademark of Nichino America, Inc.

060305

05/26/05 EPA

**NICHINO AMERICA, INC.
4550 New Linden Hill Rd., Suite 501
Wilmington, DE 19808
www.nichino.net**